arXiv:hep-ph/9402316v1 18 Feb 1994

# Quantum electrodynamics in the squeezed vacuum state: Feynman rules and corrections to the electron mass and anomalous magnetic moment

K. Svozil
Institut für Theoretische Physik
Technische Universität Wien
Wiedner Hauptstraße 8-10/136
A-1040 Vienna, Austria
e1360dab@awiuni11.edvz.univie.ac.at

February 1, 2008

## Abstract

Due to the nonvanishing average photon population of the squeezed vacuum state, finite corrections to the scattering matrix are obtained. The lowest order contribution to the electron mass shift for a one mode squeezed vacuum state is given by $\delta m(\Omega, s)/m = \alpha(2/\pi)(\Omega/m)^2 \sinh^2(s)$, where $\Omega$ and $s$ stand for the mode frequency and the squeeze parameter and $\alpha$ for the fine structure constant, respectively. The correction to the anomalous magnetic moment of the electron is $\delta a_e(s) = -(4\alpha/\pi)\sinh^2(s)$.

The dependece of the scattering matrix on the vacuum state of the theory and on exterior parameters has been studied for the thermal equilibrium [1], in cavity–quantum electrodynamics [2], on fractal space–time support [3] and, to some extent, in the presence of strong electromagnetic fields [4, 5]. Here, quantum electrodynamics is investigated in the presence of squeezed vacuum fluctuations [6]; i.e., fluctuations with reduced noise in amplitude or phase.

The squeezed vacuum state [7] exhibits a nonvanishing average photon density proportional to $\sinh^2(s)$ per squeezed mode, where $s$ is the squeeze parameter [8]. This can be accounted for in the perturbation series by the introduction of a causal photon propagator as follows [9]. Denote the squeezed vacuum by $|\mathrm{sv}\rangle$. The photon propagator in the Landau gauge is

$$D_{\mu\nu}(x-y) \quad = \quad -i\langle \mathrm{sv}|T[A_\mu(x)A_\nu(y)]|\mathrm{sv}\rangle$$

$$
\begin{aligned}
&= ig_{\mu\nu}\int \frac{d^3k}{(2\pi)^3}\frac{dk'^3}{2(E_kE_{k'})^{1/2}}\langle \mathrm{sv}|\theta(x_0-y_0)[e^{-i(kx-k'y)}a_ka^\dagger_{k'}+e^{i(kx-k'y)}a^\dagger_ka_{k'}]+x\leftrightarrow y|\mathrm{sv}\rangle \\
&= ig_{\mu\nu}\left\{\int \frac{d^3k}{(2\pi)^3}\frac{1}{2E_k}[\theta(x_0-y_0)e^{-ik(x-y)}+\theta(y_0-x_0)e^{ik(x-y)}]+\right. \\
&\qquad \left. +\int \frac{d^3k}{(2\pi)^3}\frac{1}{2E_k}n(k)[e^{ik(x-y)}+e^{-ik(x-y)}]\right\}, \qquad (1)
\end{aligned}
$$

where the $aa^\dagger$ terms generate the usual causal propagator while the $a^\dagger a$ terms count the particle density in the squeezed vacuum. The propagator can be rewritten using contour–integral techniques

$$
\begin{aligned}
D_{\mu\nu}(x-y) &= \int \frac{d^4k}{(2\pi)^4}e^{-ik(x-y)}D_{\mu\nu}(k) \\
D_{\mu\nu}(k) &= -g_{\mu\nu}\left[\frac{1}{k^2+i\epsilon}-2\pi i\delta(k^2)n(k)\right] \quad . \qquad (2)
\end{aligned}
$$

For the one mode squeezed state, $n(k;\Omega,s)=\Omega\sinh^2(s)\delta(E_k-\Omega)$, where $E_k$ is the photon energy parameter and $\Omega$ and $s$ stand for the frequency of the squeezed mode and the squeezing parameter, respectively. The electron propagator $S(p)=1/(\not p-m+i\epsilon)$, as well as the bare vertex $\gamma_\mu$ remain unchanged. Notice however that a preferred frame of reference has been introduced due to the noncovariant choice of the density $n(k;\Omega,s)$, i.e., the one at rest with respect to the squeezed vacuum. The resulting breakdown of Lorentz invariance necessitates a careful interpretation of the usual renormalisation prescriptions.

In what follows, the lowest order correction to the radiative mass of the electron will be calculated. This can be done by evaluating the second order contribution to the self energy of the electron

$$
\Sigma(p;\Omega,s)=-ie^2\int\frac{d^4k}{(2\pi)^4}[iD_{\mu\nu}(k;\Omega,s)]\gamma^\mu\frac{i}{\not p-\not k-m}\gamma^\nu \quad . \qquad (3)
$$

The physical mass is interpreted as usual as the pole of the renormalized electron propagator. For $\delta m(\Omega,s)\ll m$,

$$
\begin{aligned}
m(\Omega,s) &\approx m+\delta m+\Sigma(p;\Omega,s)|_{\not p=m} \\
&\qquad = m+\delta m+\Sigma(p;s=0)|_{\not p=m}+\delta\Sigma(p;\Omega,s)|_{\not p=m} \\
&\qquad = m+\delta m(\Omega,s) \quad , \qquad (4)
\end{aligned}
$$

where $m$ stands for the renormalized unsqueezed mass of the electron.

The correction term $\delta m(\Omega,s)=\delta\Sigma(p;\Omega,s)|_{\not p=m}$ due to squeezing adds up coherently to the renormalization contributions of $m$. Its explicit form is given by

$$
\delta m(\Omega,s) = -\frac{e^2}{(2\pi)^3}\int d^4k\delta(k^2)n(k;\Omega,s)\gamma_\mu\frac{\not p-\not k+m}{(p-k)^2-m^2+i\epsilon}\gamma^\mu\ |_{\not p=m}
$$

$$= \frac{\alpha}{2\pi^2} \frac{I_\mu(p) p^\mu}{m} |_{p^2=m^2} \quad , \tag{5}$$

where Gordon's identity which reduces to $\gamma_\mu = p_\mu/m$ has been used, $\alpha = e^2/4\pi$ stands for the fine structure constant and

$$I_\mu(p) = \int d^3\vec{\mathbf{k}} \frac{k_\mu}{|\vec{\mathbf{k}}|(pk)} n(|\vec{\mathbf{k}}|; \Omega, s) \quad . \tag{6}$$

In the rest frame of the electron this expression can be evaluated, yielding

$$\delta m(\Omega, s)/m = \alpha(2/\pi)(\Omega/m)^2 \sinh^2(s) \quad . \tag{7}$$

For optical frequencies, $\delta m(s)/m \approx 10^{-13} \sinh^2(s)$.

The correction $\delta a_e$ to the anomalous magnetic moment of the electron $a_e$ can be extracted from a decomposition of the vertex function $\Lambda_\mu = \gamma_\mu + \Gamma_\mu$ on shell

$$\bar{u}(p-q)\Lambda_\mu u(p) = \bar{u}(p-q)\left[\gamma_\mu f_1(q^2) + \frac{i}{2m}\sigma_{\mu\nu}q^\nu f_2(q^2)\right] u(p) \tag{8}$$

with $a_e = f_2(q^2 = 0)$. To lowest order one obtains

$$\begin{aligned} \Gamma_\mu(p, p-q; \Omega, s) &= \Gamma_\mu(p, p-q; s=0) + \delta\Gamma_\mu(p, p-q; \Omega, s) \\ &= (-ie)^2 \int \frac{d^4k}{(2\pi)^4} [iD_{\alpha\beta}(k; \Omega, s)] \gamma^\alpha \frac{i}{\not p - \not q - \not k - m} \gamma_\mu \frac{i}{\not p - \not k - m} \gamma^\beta \end{aligned} \tag{9}$$

The correction due to squeezing for $p = (m, 0, 0, 0)$ and $q = (q_0, 0, 0, 0)$, $q_0^2 \ll m^2$ can be written as

$$\delta\Gamma_\mu(p, p; \Omega, s) = -(i\alpha/2\pi^2)(m^2 I \gamma_\mu - I_{\mu\nu}\gamma^\nu) \tag{10}$$

$$I = \int d^4k \frac{\delta(k^2) n(k; \Omega, s)}{(pk)^2} \tag{11}$$

$$I_{\mu\nu} = \int d^4k \frac{k_\mu k_\nu \delta(k^2) n(k; \Omega, s)}{(pk)^2} \quad . \tag{12}$$

For a definition of $a_e$ the term proportional to $\sigma_{\mu\nu}q^\nu$ is chosen. Hence, only the first term proportional to $I\gamma_\mu$ on the right hand side of (10) is relevant. Using Gordon's identity, which is $\gamma_\mu = (1/2m)(2p_\mu - q_\mu - i\sigma_{\mu\nu}q^\nu)$ here, one obtains for the correction to the anomalous magnetic moment of the electron

$$\delta a_e(s) = -(4\alpha/\pi) \sinh^2(s). \tag{13}$$

This correction to the anomalous magnetic moment of the electron becomes comparable to the unsqueezed value $a_e = \alpha/2\pi$ for $s \approx 0.35$ and increases rapidly as the population of the squeezed vacuum increases and should be testable with realistic experimental parameters.

# References

[1] for a recent review see G. Barton, *Annals of Physics (N.Y.)* **200**, 271 (1990).

[2] M. Kreuzer and K. Svozil, *Phys. Rev.* **D34**, 1429 (1986); G. Barton and N. S. J. Fawcett, *Phys. Rep.* **170**, 1 (1988).

[3] A. Zeilinger and K. Svozil, *Phys. Rev. Lett.* **54**, 2553 (1985); K. Svozil and A. Zeilinger, *Journal of Modern Physics* **A1**, 971–990 (1986); K. Svozil, *J. Phys.* **A19**, L1125 (1986); *ibid.* **A20**, 3861 (1987).

[4] W. Greiner, B. Müller and J. Rafelski, *Quantum Electrodynamics of Strong Fields* (Springer, Berlin 1985).

[5] for instance *Issues in Intense–Field Quantum Electrodynamics*, ed. by V. L. Ginzburg (Nova Science Publishers, Commack, New York, 1987).

[6] for a calculation of corrections to the Lamb shift which used different methods, see G. J. Milburn, *Phys. Rev.* **A34**, 4882 (1986).

[7] R. Loudon and P. L. Knight, *Journal of Modern Optics* **34**, 709 (1987).

[8] other squeezing parameters associated with more general canonical transformations [isomorphic to elements of the real symplectic group $Sp(2N,R)$ for $N$–mode squeezing] have been set zero, but a generalisation is straightforward.

[9] In the following, units are used such that $\hbar = c = 1$. The notation of J. D. Bjorken and S. D. Drell, *Relativistic Quantum Mechanics* (McGraw–Hill, New York, 1964) is adopted.